



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

Mechanisms of Subcutaneous and Sublingual Aeroallergen Immunotherapy

Komlósi, Zsolt István ; Kovács, Nóra ; Sokolowska, Milena ; van de Veen, Willem ; Akdis, Mübeccel ;
Akdis, Cezmi A

DOI: <https://doi.org/10.1016/j.iac.2019.09.009>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-180550>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Komlósi, Zsolt István; Kovács, Nóra; Sokolowska, Milena; van de Veen, Willem; Akdis, Mübeccel; Akdis, Cezmi A (2020). Mechanisms of Subcutaneous and Sublingual Aeroallergen Immunotherapy. Immunology and Allergy Clinics of North America, 40(1):1-14.

DOI: <https://doi.org/10.1016/j.iac.2019.09.009>

Mechanisms of Subcutaneous and Sublingual Aeroallergen Immunotherapy: What Is New?



Zsolt István Komlósi, MD, PhD^{a,*}, Nóra Kovács, MD^{a,b},
Milena Sokolowska, MD, PhD^{c,d}, Willem van de Veen, PhD^{c,d},
Mübeccel Akdis, MD, PhD^{c,d}, Cezmi A. Akdis, MD^{c,d}

KEYWORDS

- Allergy • Aeroallergen • Allergen immunotherapy
- Subcutaneous allergen immunotherapy • Sublingual allergen immunotherapy
- Treg cells • Breg cells • Innate lymphoid cells

KEY POINTS

- The two most widely used forms of allergen immunotherapy (AIT) for treatment of aeroallergen-induced allergic diseases are the subcutaneous and the sublingual AIT. Substantial developments in AIT took place in the last decade to overcome problems in standardization, efficacy, safety, long duration of treatment, and costs.
- However, accurate descriptions of AIT-responsive disease endotypes with well-defined biomarkers continues to be an unmet need.
- AIT uses general mechanisms of immune tolerance to allergens with changes in allergen-specific memory T- and B-cell responses, regulation of allergen-specific IgE and IgG production, and modification of mast cell and basophil activation thresholds or dendritic cell phenotypes.
- The main highlights of recent years are advances in the understanding of innate lymphoid cells (ILC) including involvement of type 2 ILCs in development of allergic airway inflammation, and contribution of type 3 ILCs to B_{reg} cell-mediated immune tolerance.
- There is a need for further investigation of AIT mechanisms and biomarkers to identify the best candidates for AIT, the long-term responders.

^a Department of Genetics, Cell- and Immunobiology, Semmelweis University, Nagyvárad Sqr. 4, Budapest 1089, Hungary; ^b Lung Health Hospital, Munkácsy Mihály Str. 70, Törökbálint 2045, Hungary; ^c Swiss Institute of Allergy and Asthma Research (SIAR), Hermann-Burchard Strasse 9, Davos Wolfgang CH7265, Switzerland; ^d Christine Kühne – Center for Allergy Research and Education, Hermann-Burchard Strasse 1, Davos Wolfgang CH7265, Switzerland

* Corresponding author.

E-mail addresses: komlosi.zsolt@med.semmelweis-univ.hu; drkomlo@yahoo.com

INTRODUCTION

The two biggest success stories in allergy treatment are topical glucocorticosteroids and allergen immunotherapy (AIT). Glucocorticosteroids, the principal pharmaceuticals of allergy treatment, efficiently treat inflammation without curing the disease. AIT is the only therapeutic approach exerting profound effect on basic mechanisms of the disease, and is able to modify the disease course. Although glucocorticosteroid treatment also exerts some tolerance-inducing effects via its regulatory T (T_{reg}) cell-inducing capacity,^{1,2} AIT is considered to be the only treatment option with the promise of healing and induction of long-lasting tolerance, persisting even after discontinuation of therapy. Glucocorticosteroid treatment is efficient, but not effective enough, whereas AIT is effective but not efficient enough.³ Consequently, aiming for the ideal therapy to overcome the global burden of allergy, a disease-modifying treatment, such as AIT, is the best choice. Substantial progress has been made in the last decade to overcome problems in standardization, efficacy, safety, long treatment duration, patient adherence, and costs of AIT, and as a result, new guidelines have been implemented.^{4–12} Here, we summarize some major advancements of the last years in the understanding of AIT mechanisms. This review is focused on recent findings and better understanding of the mechanisms of subcutaneous and sublingual aeroallergen immunotherapy (SCIT and SLIT, respectively).

Difficulties in the management of allergic disease originate from its heterogeneity in patient population. Phenotyping allergic diseases helps identify clinical and morphologic characteristics and unique responses to treatment. However, phenotypes do not seem to be directly linked to molecular mechanisms of the disease. Endotypes of allergy and asthma describe certain mechanisms of the subgroups of the diseases, such as metabolic, inflammatory, immunologic, and remodeling pathways involved in the pathogenesis.^{13–15} The relationship of the endotypes to AIT responsiveness is unclear thus far. It is hoped that detailed description of allergy endotypes and identification of their biomarkers will enable more successful, endotype-driven AIT strategies in the future. In particular, AIT-responsive endotypes should be identified and biomarkers of these endotypes should be discovered and become available at the point of care. These biomarkers should define AIT-responsive patients as early as possible.

ROLE OF VARIOUS T CELLS IN ALLERGEN TOLERANCE AND IMMUNOTHERAPY

Regulatory T Cells

Immune tolerance to allergens implies changes in memory-type allergen-specific T- and B-cell responses, regulation of allergen-specific IgE and IgG production, and modification of mast cell and basophil activation thresholds or dendritic cell (DC) phenotypes (Fig. 1). Different levels of evidence from direct in vivo data and in vitro experiments demonstrate the role of allergen-specific T_{reg} cells in allergen tolerance in humans.

Strong in vivo evidence has been obtained by investigation of biopsies of affected tissue and skin late-phase responses. A decrease in T helper cell type 2 (T_H2) and eosinophils and a parallel increase in T_{reg} cells and their cytokines in the tissue is observed after AIT.¹⁶ In the last two decades, the pivotal role of T_{reg} cells in induction and maintenance of immune tolerance has been demonstrated. This is further supported by the fact that adoptive transfer of T_{reg} cells, depending on the timing of administration, can play a role in the prevention or treatment of several T-cell-mediated diseases, including allergic airway inflammation and hyper-responsiveness,¹⁷ a large spectrum of autoimmune diseases (eg, diabetes mellitus, experimental autoimmune encephalomyelitis, inflammatory bowel disease,




Fig. 1. Overview of the basic mechanisms of allergen immunotherapy. Regulatory T and B cells play a central role in the suppression of allergic inflammation, mainly by means of their interleukin-10 and transforming growth factor- β production. *Green arrows* indicate induction; *red arrows* indicate inhibition. B_{reg}, regulatory B cell; IL, interleukin; TGF, transforming growth factor.

systemic lupus erythematosus, rheumatoid arthritis), and graft-versus-host disease after allogeneic hematopoietic stem cell transplantation; and inhibit rejection of the graft after allogeneic solid organ transplantation in mouse models of these conditions. The human application of adoptive T_{reg} cell transfer therapy is currently being pursued in many studies.^{18,19} In the clinical setting, SCIT and SLIT have been shown to induce allergen-specific T_{reg} cells in humans. Peripheral T-cell tolerance is characterized mainly by generation of allergen-specific T_{reg} cells and decrease of T_H2 cells.^{20–24}

Direct ex vivo analysis of human peripheral blood cells without any in vitro culture also provided evidence to support the role of T_{reg} cells in maintenance of immune tolerance. Samples were collected from patients allergic to bee venom during AIT and from healthy beekeepers, who were not allergic to bee venom, on acquiring natural tolerance to high-dose allergen exposure during beekeeping season. Allergen-specific CD4⁺ T cells, identified using major histocompatibility complex class II tetramers, and interleukin (IL)-4, interferon (IFN)- γ , and IL-10 cytokine-secreting cells have been analyzed in these studies. These data demonstrate that allergen-specific T_{reg} cells increase and allergen-specific and nonspecific T_H2 cells decrease in parallel with the induction of natural or clinical immune tolerance in beekeepers during beekeeping season and AIT-treated patients with allergy, respectively.^{24,25} AIT also promotes allergen-specific IL-22 and IFN- γ producing T_H cells.^{24,26} It is now well understood that allergen-specific T_{reg} cells and allergen-specific and nonspecific effector T cells contain several distinct phenotypic compartments.²⁷ AIT involves upregulation of the activated T_{reg} and downregulation of allergen-specific immunoglobulin-like transcript 3 (ILT3)-expressing, dysfunctional T_{reg} cells.²⁴ ILT3⁺ T_{reg} cells were identified as dysfunctional T_{reg} cells, because they have impaired suppressive potency and they are unable to control the maturation of T_H2-inducing DCs.²⁸

Additional evidence has been obtained from cell cultures. IL-10 and transforming growth factor- β are produced by antigen-specific T_{reg} cells. T_{reg} cells from donors

with atopy have a reduced capability to suppress T-cell proliferation. Immune tolerance induced by SLIT is associated with an increase in T_{reg} cell numbers; elevated IL-10 production in sublingual FOXP3-expressing T_{reg} cells; and elevated allergen-specific IgG₄, IgA levels and serum blocking activity for IgE-facilitated allergen binding (IgE-FAB) to B cells.^{23,29,30} IL-10-producing T_{reg} cells suppress T_H2-type immune response and IL-17-producing T_H cells.^{31,32}

It has been recently demonstrated that a significant increase in the numbers of allergen-specific FOXP3⁺Helios⁺CD25⁺CD127⁻ T_{reg} cells took place at the end of up-dosing phase (10 weeks), which only slightly decreased at the end of the maintenance phase (3 years) of house dust mite (HDM) SCIT.²⁴ In contrast, ILT3⁺ dysfunctional T_{reg} cells²⁸ decreased substantially after 3 years of SCIT.²⁴ Additionally, IL-10⁺ allergen-specific T_{reg} cells are still present at high frequency at the end of SCIT. Increased number of FOXP3⁺Helios⁺ and IL-10⁺ T_{reg} cell and decreased ILT3⁺ T_{reg} cell responses correlated with improved allergic symptoms. These data further confirm that the induction of allergen-specific functional T_{reg} cells is one of the key events of AIT.²⁴ In all of these mechanisms, the H₂ receptor plays an important role. It has been shown that it can suppress DCs, T cells, natural killer T cells, and basophils. This results in antagonizing the proallergic effects of histamine via the H₁ receptor.^{33–35} Histamine signaling is a newly determined link between microbiome and immune tolerance. It has been recently shown that histamine-secreting commensal bacteria exist in human gut and are linked to severity of asthma³⁶ and have pleiotropic effects on tolerance in the lungs.^{37,38}

Taken together, the critical role for allergen-specific T_{reg} cell-mediated immune tolerance in successful AIT is well established. However, this is not the only mechanism operational in the AIT-induced tolerance.

The equally important immunologic phenomenon that warrants the favorable clinical outcome of AIT is the eradication of allergen-specific T_H2 cells (Fig. 2). It has been recently demonstrated that there are subsets of allergen-specific T_H2 cells, such as CD27⁻CD45RB⁻CRTH2⁺CD49d⁺CD161⁺ T_H2 memory cells, CRTH2⁺CCR4⁺CD27⁻CD4⁺ T_H2 cells,³² or ST2⁺CD45RO⁺CD4⁺ cells,³⁹ of special importance in the pathogenesis of allergic diseases. These so-called T_H2A cells or pathogenic T_H2 cells are specifically targeted during AIT, and their eradication is indicative of favorable clinical response.^{3,39,40} These cells are characterized by high




Fig. 2. Direction of change in some important T-cell subsets in response to AIT. Allergen-specific pathogenic T cells decrease and regulatory T cells increase during a successful AIT.

expression of *IL17RB* (gene encoding IL-25 receptor), *IL1RL1* (gene encoding ST2, a receptor for IL-33), and *CRLF2* (gene encoding TSLP receptor), and they highly express *IL13* and *IL5* genes.^{40,41}

Interleukin-35

IL-35 is a newly identified inhibitory cytokine produced by T_{reg} cells and is essential for their immunosuppressive function.⁴² IL-35-inducible regulatory T (iT_{R35}) cell is a newly recognized subset of induced T_{reg} cells with IL-35⁺CD4⁺CD25⁺FOXP3⁻ phenotype and potent immune regulatory properties.⁴³ The role of IL-35 and IL-35-producing iT_{R35} cells in SLIT has recently been described (see Fig. 2). Proliferation and cytokine production of allergen-specific T_H2 effector cells, IL-25- or IL-33-induced IL-5 and IL-13 production of innate lymphoid cells (ILC) 2 cells, and CD40L-, IL-4-, and IL-21-mediated IgE production of B cells can all be inhibited by IL-35, whereas the production of IFN- γ and IL-10 is induced by this cytokine. Moreover, iT_{R35} cells reduce memory T-cell proliferation and T_H2 cytokine production in an IL-10- and IL-35-dependent manner. Of note, the number of iT_{R35} cells and allergen-induced IL-35 production is higher in the peripheral blood of patients with allergic rhinitis after successful SLIT than in the blood of untreated patients, and similar to that of healthy control subjects. Both the percentage and the IL-35-producing capacity of the iT_{R35} cells inversely correlate with rhinitis symptom score.⁴⁴ Taken together, IL-35 and its source (iT_{R35} cell) may represent a new target for biomarker development and for future therapies.⁴⁵

Follicular Helper T Cells

Increasing evidence indicates that the regulation of follicular helper T (T_{FH}) cells is also of great importance in the success of AIT (see Fig. 2). Circulating T_{FH} cells are divided into subgroups corresponding to T_{H1} , T_{H2} , and T_{H17} helper cells. IgE antibody production by B cells is supported by type 2 T_{FH} (T_{FH2}) cells.⁴⁶ T_{FH2} cell numbers are increased in the peripheral blood of patients with allergic rhinitis, and are able to induce IgE production on an IL-4-dependent manner. Moreover, allergen-specific IgE levels correlate only with allergen-specific T_{FH2} cell frequencies, but not with non-follicular T_H2 cells frequencies in patients allergic to HDM. The numbers of T_{FH2} cells reduced more sharply than T_H2 cells in response to AIT. Of note, the combined symptom and medication score improvement in response to SCIT correlated only with the percent reduction of allergen-specific T_{FH2} cell numbers, and not with that of T_H2 cells.⁴⁷ T_{FH} activities are modulated by follicular T_{reg} (T_{FR}) cells⁴⁸ and by regulatory B (B_{reg}) cells.⁴⁹ T_{FR} cells were first described in human tonsils and differentiate from natural T_{reg} cells.⁵⁰ T_{FR} cell frequencies are decreased in tonsils of patients allergic to HDM suffering from allergic rhinitis. Tonsillar and blood T_{FR} cell phenotypes and numbers correlate with each other. Circulating T_{FR} cells of patients with allergic rhinitis fail to suppress IgE production of B cells in vitro. Circulating T_{FR} cell frequencies are increased and their IgE-suppressing function is improved after AIT.⁵¹ Therefore, therapeutic targeting of T_{FR} and T_{FH} cells and their use as a biomarker has been proposed.

INNATE LYMPHOID CELLS IN ALLERGY AND ALLERGEN IMMUNOTHERAPY

Exploration of the diversity of ILCs has substantially changed the understanding of tissue inflammation and immune tolerance in the last few years.⁵² ILCs are developmentally related to natural killer cells. Novel members of the ILC family (ILC1s, ILC2s, and ILC3s) show similar biologic characteristics, including transcription factor expression and cytokine production, to the key cellular elements of adaptive immune response,

the T_H cell subpopulations: T_H1, T_H2, and T_H17/T_H22, respectively.⁵³ The whole genome signatures of different ILCs subpopulations and ILC heterogeneity in humans have recently been published.^{54,55} Particularly, ILC2s play an important role in allergic airway inflammation.⁵⁶ They are dependent on GATA3, and they substantially contribute to IL-5 and IL-13 production in allergic airway responses.⁵⁷ ILC3s are ROR γ t-dependent cells. Their signature-cytokine is IL-22, an important mediator of the mucosal immune defense, tissue repair, and the maintenance of epithelial barrier integrity.⁵⁸

It has recently been demonstrated that seasonal increase in peripheral ILC2s is inhibited by grass pollen extract SCIT.⁵⁹ Cross-sectional studies suggest that the relative proportion of circulating ILCs is affected by AIT. The ILC2/ILC1 ratio is increased in patients allergic to HDM suffering from allergic rhinitis⁶⁰; however, it is normalized by a successful AIT. Nonresponders show similar ILC2/ILC1 ratio to untreated patients with allergic rhinitis.^{61,62}

Our group demonstrated that IL-10⁺ regulatory ILCs (ILC_{reg}s) are induced from ILC2s by retinoic acid.⁶³ CCR2 and ST2 (hallmarks of ILC2s) are downregulated and CD25 and CTLA-4 are upregulated on ILC_{reg}s. These cells are able to suppress T-cell and ILC2 activation. Retinoic acid-producing DCs were found to help peripheral T_{reg} cell differentiation.^{64,65} Retinoic acid-induced regulatory DCs have a protective effect on adoptive transfer in a mouse model of food-induced anaphylaxis.⁶⁶ Considering the general role of retinoic acid in tolerance-induction, the participation of ILC_{reg}s in the mechanisms of AIT-induced tolerance is also possible.

Moreover, we demonstrated recently that activated, CD40L-expressing ILC3s reside on the border of the T cell–B cell areas in tonsils, and are in close contact with B cells *in vivo*. Furthermore, we showed that CD40L⁺ ILC3s and B cells are in a mutually beneficial relationship with each other: ILC3s induce IL-15 production in B cells via BAFF-receptor, whereas IL-15, a potent growth factor for ILC3s, upregulates CD40L expression on ILC3s. IL-15-activated CD40L⁺ ILC3s help B-cell survival, proliferation and differentiation of IL-10 secreting, and functional B_{reg} cells in a CD40L- and BAFF-receptor-dependent manner. Moreover, ILC3-induced B_{reg} cells dominantly have CD27-IgD⁺IgM⁺CD24^{high} CD38^{high} CD1d⁺ immature transitional (itB_{reg}) phenotype. This mechanism contributes to the maintenance of immune tolerance to innocuous antigens and becomes insufficient in allergic diseases.⁶⁷ Tonsils are important mucosal sites of immune tolerance,⁶⁸ where generation of functional allergen-specific T_{reg} cells occurs. We showed that ILC3s and B_{reg} cells colocalize in the interfollicular regions of palatine tonsils, together with T_{reg} cells. These data suggest that there are regulatory niches in tonsils where T_{reg} and B_{reg} cells develop next to each other. CD40L⁺ ILC3s might be involved in maintenance of immune tolerance in tonsils through induction of functional itB_{reg} cells. These cells can contribute to suppression of T-cell responses through cell-cell contact via programmed cell death-ligand 1 and via secreted IL-10.⁶⁷ These mechanisms can also play a role in immune tolerance induction during successful SLIT (see Fig. 1).^{35,69}

The significant reduction in total and allergen-specific IgE after 3 month of HDM SLIT was not affected by previous tonsillectomy; however, the mitigation of nasal symptoms and wheeze was less pronounced in the tonsillectomy group.⁷⁰ In fact, only the palatine tonsils and/or adenoids are removed or reduced during those operations, and the lingual tonsils remained untouched.⁶⁹ These data suggest that reduction of the lymphoid tissue mass in Waldeyer ring does not impede systemic tolerance induction by SLIT; however, it may delay the improvement of clinical symptoms. The small-scale study did not reveal reasons for this dichotomy.⁷⁰ Neither the cellular infiltration of the affected tissues by T_H2 cells and eosinophils, nor the local IgE levels (entropy) were examined, both of which certainly has an impact on the symptoms and may have been perpetuated in

tonsillectomized patients after SLIT. Larger scale, randomized, controlled studies with appropriate blinded design are needed to be able to appreciate the influence of tonsillectomy on AIT efficacy. In addition, it should be considered that lingual tonsil, which has the same structure and cellular composition as palatine tonsils, is a large lymphoid organ that remains intact in tonsillectomy patients. IgE-mediated allergy was found to be associated with substantially lower risk of complicated appendicitis in a cohort of children who underwent surgery for acute appendicitis.⁷¹ However, the influence of appendectomy on development of allergy, and specifically on the response to immunotherapy, has not been investigated thus far.

REGULATORY B CELLS AND ANTIBODY PRODUCTION IN ALLERGEN TOLERANCE AND IMMUNOTHERAPY

Regulatory B Cells

Deficiency of IL-10-producing B_{reg} cells leads to increased inflammation in mouse models of autoimmunity, transplantation, and chronic infection. Absolute numbers of T_{reg} cells and expression levels of FOXP3 in these cells are dependent on IL-10 production by B cells. Modulation of T_{reg} cell frequencies in vivo is exclusively restricted to transitional 2 marginal zone precursor B_{reg} cells in mice.⁷² Of note, iT_{Breg} cells can prevent and reverse airway inflammation in allergized mice by inducing recruitment of T_{reg} cells to the lung.⁷³ Adoptive transfer of iT_{Breg} cells can ameliorate allergic airway inflammation and hyperresponsiveness in a mouse model of allergic asthma in an IL-10-dependent manner.⁷⁴ Human iT_{Breg} cells are also able to induce T_{reg} cells in vitro.⁷⁵ The relative proportion of B_{reg} cell subsets within B cells was found to be reduced in patients with allergic diseases, such as allergic rhinitis⁷⁶ and allergic asthma,⁷⁷ and the frequency of IL-10-producing, antigen-specific B cells increased after AIT.^{78,79} In vivo evidence for allergen tolerance has been demonstrated: the suppressive B cells and IgG₄-expressing B cells are developed from IL-10⁺ BR1 cells in human subjects after AIT.⁸⁰ Supporting the role of IL-10, solely IL-10-overexpressing B cells acquired an immunoregulatory profile comprising upregulation of suppressor of cytokine signaling 3 (SOCS3), glycoprotein A repetitions predominant (GARP), CD25, and programmed cell death-ligand 1.⁸¹ These cells showed a significant reduction in levels of many proinflammatory cytokines, and augmented the production of anti-inflammatory IL-1 receptor antagonist and vascular endothelial growth factor. In addition, IL-10-overexpressing B cells secreted less IgE and showed a general T-cell and DC suppression.⁸¹ Moreover, a novel mechanism of AIT-induced specific immune tolerance is suggested by our recent data. Increased numbers of IgA- and IgG₄-expressing allergen-specific memory B cells, plasmablasts, and IL-10⁺ and/or IL-1RA⁺ B_{reg} cells were found in the peripheral blood over a 2-year period during successful HDM SCIT (see [Fig. 1](#)).⁸²

Modulation of Allergen-Specific IgE and IgG Responses During Allergen Immunotherapy

Specific IgE increases in individuals allergic to pollen during the pollen season and a successful AIT can prevent this increase. AIT itself also induces an early increase in IgE during the up-dosing phase, which rings out in maintenance phase in parallel with the increase in IgA and IgG₄ (see [Fig. 1](#)).⁸³

AIT-induced IgG may be directed against the same epitopes as IgE, resulting in direct competition for allergen binding and a blocking effect.⁸⁴ Allergen-specific IgG antibodies can inhibit IgE-mediated basophil and mast cell activation and degranulation with two different mechanisms: allergen neutralization and binding to the inhibitory FcγRIIb. It has been demonstrated recently that although only high-affinity IgG

antibodies can have both of these effects, low-affinity IgG antibodies are still able to inhibit mast cell activation via Fc γ RIIb.⁸⁵ IgG₄ antibodies have several features that may play a role in their anti-inflammatory effects. Two arms of IgG₄ have the ability to separate and repair by dynamic Fab arm exchange, which leads to bispecific antibodies that are functionally monomeric.⁸⁶ Furthermore, IgG₄ does not fix complement and is capable of inhibiting immune-complex formation by other isotypes.⁸⁷ It is likely the decrease in IgE/IgG₄ ratio during AIT is a biomarker of skew from allergen-specific T_H2 to T_{reg} cell and B_{reg} cell predominance. IL-10 is a potent suppressor of total and allergen-specific IgE, whereas it simultaneously increases IgG₄ production.^{21,88}

The characteristic changes in the level of IgE and IgG₄ during SCIT is mirrored by the local concentrations of those immunoglobulins in nasal secretions⁴⁵ and saliva.⁸⁹

The results of some recent studies confirm that the IgG₄-mediated inhibition of CD23-dependent IgE-FAB,⁹⁰ rather than the serum concentration of IgG₄ solely, is a valuable biomarker of the AIT clinical outcome.^{29,91} Other investigators evidenced that specific IgG₄ and inhibition of IgE-FAB correlates well with clinical efficacy of AIT.⁹² Inhibition of IgE-FAB by nasal secretions correlates more closely with clinical outcome than that of serum samples after SCIT.⁴⁵

Regulation of Mast Cells and Basophils by Allergen Immunotherapy

Clinical desensitization starts as early as after the first injection of SCIT. This is mainly linked to decreased mast cell and basophil degranulation, which has been demonstrated in a mouse model.⁹³ One of the main soluble factors released from mast cells and basophils following allergen challenge is histamine, which mediates its effects via histamine receptors. Immunosilencing of Fc ϵ RI-activated basophils by means of selective suppression mediated by H₂ receptor might be highly relevant for the early desensitization effect, at least in venom AIT.⁹⁴ Although there are individual differences and risks for developing systemic anaphylaxis, the suppression of mast cells and basophils is further increased during the course of AIT. Early reduction in basophil sensitivity predicts symptom relief also in grass pollen immunotherapy.⁹⁵ Furthermore, basophil expression of diamine oxidase shows a significant increase after AIT and is suggested as a novel biomarker of AIT responsiveness.⁹⁶

OTHER ASPECTS OF IMMUNOTHERAPY

It has been demonstrated in mouse model of allergic asthma in vivo and human bronchial epithelial cell cultures in vitro that AIT contributes to the restoration of the airway epithelial barrier integrity by the reduction of IL-25 production. IL-25 is one of the main inducers of endoplasmic reticulum stress and tight junction damage.⁹⁷ It has been recently shown in a 5-year clinical study that grass pollen SLIT significantly reduces the risk of experiencing asthma symptoms and decreases the need for asthma medications.⁹⁸ The systematic meta-analyses of contribution of AIT in allergic asthma⁹⁹ and allergic rhinoconjunctivitis¹⁰⁰ have been recently published, supporting the previously mentioned study and confirming that AIT has positive effects on various allergic diseases. The effects of AIT for various phenotypes and endotypes of allergic diseases need to be further elucidated in vivo in animal models and directly in human tissue.^{101,102}

SUMMARY

AIT offers an efficient treatment of allergic diseases with a possibility of cure. There is strong evidence for clinical safety and efficacy of AIT. Demonstration of allergen-specific T- and B-cell tolerance, mediated, among others, by the immunosuppressive

functions of IL-10, led to a major conceptual change in this area and opened a way to bring AIT from empiricism to a treatment with solid mechanisms of action linked to immune tolerance. The accurate definition of disease endotypes and a correct selection of the responder patients with well-defined biomarkers still remain essential unmet needs in the clinical setting. Therefore, deeper understanding of immuno-pharmaco-metabolo-genomics of allergy and the impact of the environmental exposure and the microbiome on it will probably help improve the AIT to accomplish vaccination strategies that are suitable for whole population-based allergy prevention and cure in the future.

ACKNOWLEDGMENTS

The laboratory of the corresponding author Z.I. Komlósi is supported by Hungarian Pediatric Oncology Network. The laboratories of the authors C.A. Akdis and M. Akdis are supported by the Swiss National Science Foundation.

REFERENCES

1. Hamdi H, Godot V, Maillot MC, et al. Induction of antigen-specific regulatory T lymphocytes by human dendritic cells expressing the glucocorticoid-induced leucine zipper. *Blood* 2007;110(1):211–9.
2. Karaki S, Garcia G, Tcherakian C, et al. Enhanced glucocorticoid-induced leucine zipper in dendritic cells induces allergen-specific regulatory CD4(+) T-cells in respiratory allergies. *Allergy* 2014;69(5):624–31.
3. Renand A, Shamji MH, Harris KM, et al. Synchronous immune alterations mirror clinical response during allergen immunotherapy. *J Allergy Clin Immunol* 2018; 141(5):1750–60.e1.
4. Halken S, Larenas-Linnemann D, Roberts G, et al. EAACI guidelines on allergen immunotherapy: prevention of allergy. *Pediatr Allergy Immunol* 2017;28(8): 728–45.
5. Bonertz A, Roberts GC, Hoefnagel M, et al. Challenges in the implementation of EAACI guidelines on allergen immunotherapy: a global perspective on the regulation of allergen products. *Allergy* 2018;73(1):64–76.
6. Larenas-Linnemann DES, Antolin-Amerigo D, Parisi C, et al. National clinical practice guidelines for allergen immunotherapy: an international assessment applying AGREE-II. *Allergy* 2018;73(3):664–72.
7. Roberts G, Pfaar O, Akdis CA, et al. EAACI guidelines on allergen immunotherapy: allergic rhinoconjunctivitis. *Allergy* 2018;73(4):765–98.
8. Bonertz A, Roberts G, Slater JE, et al. Allergen manufacturing and quality aspects for allergen immunotherapy in Europe and the United States: an analysis from the EAACI AIT guidelines project. *Allergy* 2018;73(4):816–26.
9. Ryan D, Gerth van Wijk R, Angier E, et al. Challenges in the implementation of the EAACI AIT guidelines: a situational analysis of current provision of allergen immunotherapy. *Allergy* 2018;73(4):827–36.
10. Sturm GJ, Varga EM, Roberts G, et al. EAACI guidelines on allergen immunotherapy: hymenoptera venom allergy. *Allergy* 2018;73(4):744–64.
11. Pajno GB, Fernandez-Rivas M, Arasi S, et al. EAACI Guidelines on allergen immunotherapy: IgE-mediated food allergy. *Allergy* 2018;73(4):799–815.
12. Agache I, Lau S, Akdis CA, et al. EAACI guidelines on allergen immunotherapy: house dust mite-driven allergic asthma. *Allergy* 2019;74(5):855–73.
13. Akdis CA, Bachert C, Cingi C, et al. Endotypes and phenotypes of chronic rhinosinusitis: a PRACTALL document of the European Academy of Allergy and

- Clinical Immunology and the American Academy of Allergy, Asthma & Immunology. *J Allergy Clin Immunol* 2013;131(6):1479–90.
14. Wenzel SE. Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat Med* 2012;18(5):716–25.
 15. Sugita K, Sokolowska M, Akdis CA. Key points for moving the endotypes field forward. In: Agache I, Hellings P, editors. Implementing precision medicine in best practices of chronic airway diseases. London: Elsevier; 2018. p. 107–14.
 16. Radulovic S, Jacobson MR, Durham SR, et al. Grass pollen immunotherapy induces Foxp3-expressing CD4⁺ CD25⁺ cells in the nasal mucosa. *J Allergy Clin Immunol* 2008;121(6):1467–72, 1472.e1.
 17. Kearley J, Barker JE, Robinson DS, et al. Resolution of airway inflammation and hyperreactivity after in vivo transfer of CD4⁺CD25⁺ regulatory T cells is interleukin 10 dependent. *J Exp Med* 2005;202(11):1539–47.
 18. Romano M, Fanelli G, Albany CJ, et al. Past, present, and future of regulatory T cell therapy in transplantation and autoimmunity. *Front Immunol* 2019;10:43.
 19. Trzonkowski P, Bacchetta R, Battaglia M, et al. Hurdles in therapy with regulatory T cells. *Sci Transl Med* 2015;7(304):304ps318.
 20. Akdis CA, Akdis M, Blesken T, et al. Epitope specific T cell tolerance to phospholipase A₂ in bee venom immunotherapy and recovery by IL-2 and IL-15 *in vitro*. *J Clin Invest* 1996;98:1676–83.
 21. Akdis CA, Blesken T, Akdis M, et al. Role of interleukin 10 in specific immunotherapy. *J Clin Invest* 1998;102(1):98–106.
 22. Akdis CA, Akdis M. Mechanisms of immune tolerance to allergens: role of IL-10 and Tregs. *J Clin Invest* 2014;124(11):4678–80.
 23. Suarez-Fueyo A, Ramos T, Galan A, et al. Grass tablet sublingual immunotherapy downregulates the TH2 cytokine response followed by regulatory T-cell generation. *J Allergy Clin Immunol* 2014;133(1):130–8.e1-2.
 24. Boonpiyathad T, Sokolowska M, Morita H, et al. Der p 1-specific regulatory T-cell response during house dust mite allergen immunotherapy. *Allergy* 2019;74(5):976–85.
 25. Meiler F, Zumkehr J, Klunker S, et al. In vivo switch to IL-10-secreting T regulatory cells in high dose allergen exposure. *J Exp Med* 2008;205(12):2887–98.
 26. van de Veen W, Wirz OF, Globinska A, et al. Novel mechanisms in immune tolerance to allergens during natural allergen exposure and allergen-specific immunotherapy. *Curr Opin Immunol* 2017;48:74–81.
 27. Zemmour D, Zilionis R, Kiner E, et al. Publisher correction: single-cell gene expression reveals a landscape of regulatory T cell phenotypes shaped by the TCR. *Nat Immunol* 2018;19(6):645.
 28. Ulges A, Klein M, Reuter S, et al. Protein kinase CK2 enables regulatory T cells to suppress excessive TH2 responses in vivo. *Nat Immunol* 2015;16(3):267–75.
 29. Scadding GW, Shamji MH, Jacobson MR, et al. Sublingual grass pollen immunotherapy is associated with increases in sublingual Foxp3-expressing cells and elevated allergen-specific immunoglobulin G4, immunoglobulin A and serum inhibitory activity for immunoglobulin E-facilitated allergen binding to B cells. *Clin Exp Allergy* 2010;40(4):598–606.
 30. Jutel M, Akdis M, Budak F, et al. IL-10 and TGF-beta cooperate in the regulatory T cell response to mucosal allergens in normal immunity and specific immunotherapy. *Eur J Immunol* 2003;33(5):1205–14.
 31. Berings M, Karaaslan C, Altunbulakli C, et al. Advances and highlights in allergen immunotherapy: on the way to sustained clinical and immunologic tolerance. *J Allergy Clin Immunol* 2017;140(5):1250–67.

32. Scadding GW, Calderon MA, Shamji MH, et al. Effect of 2 years of treatment with sublingual grass pollen immunotherapy on nasal response to allergen challenge at 3 years among patients with moderate to severe seasonal allergic rhinitis: the GRASS randomized clinical trial. *JAMA* 2017;317(6):615–25.
33. Ferstl R, Frei R, Barcik W, et al. Histamine receptor 2 modifies iNKT cell activity within the inflamed lung. *Allergy* 2017;72(12):1925–35.
34. Ferstl R, Frei R, Schiavi E, et al. Histamine receptor 2 is a key influence in immune responses to intestinal histamine-secreting microbes. *J Allergy Clin Immunol* 2014;134(3):744–6.e3.
35. Palomares O, Martin-Fontecha M, Lauener R, et al. Regulatory T cells and immune regulation of allergic diseases: roles of IL-10 and TGF- β . *Genes Immun* 2014;15(8):511–20.
36. Barcik W, Pugin B, Westermann P, et al. Histamine-secreting microbes are increased in the gut of adult asthma patients. *J Allergy Clin Immunol* 2016;138(5):1491–4.e7.
37. Lunjani N, Satitsuksanoa P, Lukasik Z, et al. Recent developments and highlights in mechanisms of allergic diseases: microbiome. *Allergy* 2018;73(12):2314–27.
38. Barcik W, Pugin B, Bresco MS, et al. Bacterial secretion of histamine within the gut influences immune responses within the lung. *Allergy* 2019;74(5):899–909.
39. Ihara F, Sakurai D, Yonekura S, et al. Identification of specifically reduced Th2 cell subsets in allergic rhinitis patients after sublingual immunotherapy. *Allergy* 2018;73(9):1823–32.
40. Wambre E, Bajzik V, DeLong JH, et al. A phenotypically and functionally distinct human TH2 cell subpopulation is associated with allergic disorders. *Sci Transl Med* 2017;9(401) [pii:eaam9171].
41. Chiang D, Chen X, Jones SM, et al. Single-cell profiling of peanut-responsive T cells in patients with peanut allergy reveals heterogeneous effector TH2 subsets. *J Allergy Clin Immunol* 2018;141(6):2107–20.
42. Collison LW, Workman CJ, Kuo TT, et al. The inhibitory cytokine IL-35 contributes to regulatory T-cell function. *Nature* 2007;450(7169):566–9.
43. Collison LW, Chaturvedi V, Henderson AL, et al. IL-35-mediated induction of a potent regulatory T cell population. *Nat Immunol* 2010;11(12):1093–101.
44. Shamji MH, Layhadi JA, Achkova D, et al. Role of IL-35 in sublingual allergen immunotherapy. *J Allergy Clin Immunol* 2019;143(3):1131–42.e4.
45. Layhadi JA, Eguiluz-Gracia I, Shamji MH. Role of IL-35 in sublingual allergen immunotherapy. *Curr Opin Allergy Clin Immunol* 2019;19(1):12–7.
46. Morita R, Schmitt N, Bentebibel SE, et al. Human blood CXCR5(+)CD4(+) T cells are counterparts of T follicular cells and contain specific subsets that differentially support antibody secretion. *Immunity* 2011;34(1):108–21.
47. Yao Y, Chen CL, Wang N, et al. Correlation of allergen-specific T follicular helper cell counts with specific IgE levels and efficacy of allergen immunotherapy. *J Allergy Clin Immunol* 2018;142(1):321–4.e10.
48. Lim HW, Hillsamer P, Kim CH. Regulatory T cells can migrate to follicles upon T cell activation and suppress GC-Th cells and GC-Th cell-driven B cell responses. *J Clin Invest* 2004;114(11):1640–9.
49. Achour A, Simon Q, Mohr A, et al. Human regulatory B cells control the TFH cell response. *J Allergy Clin Immunol* 2017;140(1):215–22.
50. Varricchi G, Harker J, Borriello F, et al. T follicular helper (Tfh) cells in normal immune responses and in allergic disorders. *Allergy* 2016;71(8):1086–94.

51. Yao Y, Wang ZC, Wang N, et al. Allergen immunotherapy improves defective follicular regulatory T cells in patients with allergic rhinitis. *J Allergy Clin Immunol* 2019;144(1):118–28.
52. Kortekaas Krohn I, Shikhagaie MM, Golebski K, et al. Emerging roles of innate lymphoid cells in inflammatory diseases: clinical implications. *Allergy* 2018; 73(4):837–50.
53. Spits H, Artis D, Colonna M, et al. Innate lymphoid cells: a proposal for uniform nomenclature. *Nat Rev Immunol* 2013;13(2):145–9.
54. Bjorklund AK, Forkel M, Picelli S, et al. The heterogeneity of human CD127(+) innate lymphoid cells revealed by single-cell RNA sequencing. *Nat Immunol* 2016;17(4):451–60.
55. Li S, Morita H, Sokolowska M, et al. Gene expression signatures of circulating human type 1, 2, and 3 innate lymphoid cells. *J Allergy Clin Immunol* 2019;143(6): 2321–5.
56. Bartemes KR, Kephart GM, Fox SJ, et al. Enhanced innate type 2 immune response in peripheral blood from patients with asthma. *J Allergy Clin Immunol* 2014;134(3):671–8.e4.
57. Mjosberg JM, Trifari S, Crellin NK, et al. Human IL-25- and IL-33-responsive type 2 innate lymphoid cells are defined by expression of CCR2 and CD161. *Nat Immunol* 2011;12(11):1055–62.
58. Klose CS, Artis D. Innate lymphoid cells as regulators of immunity, inflammation and tissue homeostasis. *Nat Immunol* 2016;17(7):765–74.
59. Lao-Araya M, Steveling E, Scadding GW, et al. Seasonal increases in peripheral innate lymphoid type 2 cells are inhibited by subcutaneous grass pollen immunotherapy. *J Allergy Clin Immunol* 2014;134(5):1193–5.e4.
60. Zhong H, Fan XL, Yu QN, et al. Increased innate type 2 immune response in house dust mite-allergic patients with allergic rhinitis. *Clin Immunol* 2017;183: 293–9.
61. Fan DC, Wang XD, Wang CS, et al. Suppression of immunotherapy on group 2 innate lymphoid cells in allergic rhinitis. *Chin Med J (Engl)* 2016;129(23):2824–8.
62. Mitthamsiri W, Pradubpongsa P, Sangasapaviliya A, et al. Decreased CCR2 expression and response to allergen re-stimulation on innate lymphoid cells in patients with allergen-specific immunotherapy. *Allergy Asthma Immunol Res* 2018;10(6):662–74.
63. Morita H, Kubo T, Ruckert B, et al. Induction of human regulatory innate lymphoid cells from group 2 innate lymphoid cells by retinoic acid. *J Allergy Clin Immunol* 2019;143(6):2190–201.e9.
64. Coombes JL, Siddiqui KR, Arancibia-Carcamo CV, et al. A functionally specialized population of mucosal CD103+ DCs induces Foxp3+ regulatory T cells via a TGF-beta and retinoic acid-dependent mechanism. *J Exp Med* 2007;204(8): 1757–64.
65. Sun CM, Hall JA, Blank RB, et al. Small intestine lamina propria dendritic cells promote de novo generation of Foxp3 T reg cells via retinoic acid. *J Exp Med* 2007;204(8):1775–85.
66. Dawicki W, Li C, Town J, et al. Therapeutic reversal of food allergen sensitivity by mature retinoic acid-differentiated dendritic cell induction of LAG3(+)CD49b(-) Foxp3(-) regulatory T cells. *J Allergy Clin Immunol* 2017;139(5):1608–20.e3.
67. Komlósi ZI, Kovacs N, van de Veen W, et al. Human CD40 ligand-expressing type 3 innate lymphoid cells induce IL-10-producing immature transitional regulatory B cells. *J Allergy Clin Immunol* 2018;142(1):178–94.e11.

68. Palomares O, Ruckert B, Jartti T, et al. Induction and maintenance of allergen-specific FOXP3+ Treg cells in human tonsils as potential first-line organs of oral tolerance. *J Allergy Clin Immunol* 2012;129(2):510–20, 520.e1-9.
69. Moingeon P. Update on immune mechanisms associated with sublingual immunotherapy: practical implications for the clinician. *J Allergy Clin Immunol Pract* 2013;1(3):228–41.
70. Refaat M, Ashour ZA, Farres MN, et al. Effect of tonsillectomy on the efficacy of house dust mite sublingual immunotherapy. *Allergol Immunopathol (Madr)* 2015;43(1):108–11.
71. Salo M, Gudjonsdottir J, Omling E, et al. Association of IgE-mediated allergy with risk of complicated appendicitis in a pediatric population. *JAMA Pediatr* 2018;172(10):943–8.
72. Carter NA, Vasconcellos R, Rosser EC, et al. Mice lacking endogenous IL-10-producing regulatory B cells develop exacerbated disease and present with an increased frequency of Th1/Th17 but a decrease in regulatory T cells. *J Immunol* 2011;186(10):5569–79.
73. Amu S, Saunders SP, Kronenberg M, et al. Regulatory B cells prevent and reverse allergic airway inflammation via FoxP3-positive T regulatory cells in a murine model. *J Allergy Clin Immunol* 2010;125(5):1114–24.e8.
74. Braza F, Chesne J, Durand M, et al. A regulatory CD9(+) B-cell subset inhibits HDM-induced allergic airway inflammation. *Allergy* 2015;70(11):1421–31.
75. Lemoine S, Morva A, Youinou P, et al. Human T cells induce their own regulation through activation of B cells. *J Autoimmun* 2011;36(3–4):228–38.
76. Kim AS, Doherty TA, Karta MR, et al. Regulatory B cells and T follicular helper cells are reduced in allergic rhinitis. *J Allergy Clin Immunol* 2016;138(4):1192–5.e5.
77. Kamekura R, Shigehara K, Miyajima S, et al. Alteration of circulating type 2 follicular helper T cells and regulatory B cells underlies the comorbid association of allergic rhinitis with bronchial asthma. *Clin Immunol* 2015;158(2):204–11.
78. Boonpiyathad T, Meyer N, Moniuszko M, et al. High-dose bee venom exposure induces similar tolerogenic B-cell responses in allergic patients and healthy beekeepers. *Allergy* 2017;72(3):407–15.
79. van de Veen W. The role of regulatory B cells in allergen immunotherapy. *Curr Opin Allergy Clin Immunol* 2017;17(6):447–52.
80. van de Veen W, Stanic B, Yaman G, et al. IgG4 production is confined to human IL-10-producing regulatory B cells that suppress antigen-specific immune responses. *J Allergy Clin Immunol* 2013;131(4):1204–12.
81. Stanic B, van de Veen W, Wirz OF, et al. IL-10-overexpressing B cells regulate innate and adaptive immune responses. *J Allergy Clin Immunol* 2015;135(3):771–80.e8.
82. Boonpiyathad T, van de Veen W, Wirz O, et al. Role of Der p 1-specific B cells in immune tolerance during 2 years of house dust mite-specific immunotherapy. *J Allergy Clin Immunol* 2019;143(3):1077–86.e10.
83. Jimenez-Saiz R, Patil SU. The multifaceted B cell response in allergen immunotherapy. *Curr Allergy Asthma Rep* 2018;18(12):66.
84. Klunker S, Saggat LR, Seyfert-Margolis V, et al. Combination treatment with omalizumab and rush immunotherapy for ragweed-induced allergic rhinitis: inhibition of IgE-facilitated allergen binding. *J Allergy Clin Immunol* 2007;120(3):688–95.

85. Zha L, Leoratti FMS, He L, et al. An unexpected protective role of low-affinity allergen-specific IgG through the inhibitory receptor FcγRIIb. *J Allergy Clin Immunol* 2018;142(5):1529–36.e6.
86. van der Neut Kolfschoten M, Schuurman J, Losen M, et al. Anti-inflammatory activity of human IgG4 antibodies by dynamic Fab arm exchange. *Science* 2007;317(5844):1554–7.
87. Aalberse RC, Stapel SO, Schuurman J, et al. Immunoglobulin G4: an odd antibody. *Clin Exp Allergy* 2009;39(4):469–77.
88. Meiler F, Klunker S, Zimmermann M, et al. Distinct regulation of IgE, IgG4 and IgA by T regulatory cells and toll-like receptors. *Allergy* 2008;63(11):1455–63.
89. Liu Y, Xing Z, Wang J, et al. Salivary immunoglobulin A, E, and G4 levels specific to dermatophagoides pteronyssinus in allergic rhinitis patients treated with subcutaneous immunotherapy. *Am J Rhinol Allergy* 2018;32(6):458–64.
90. Shamji MH, Wilcock LK, Wachholz PA, et al. The IgE-facilitated allergen binding (FAB) assay: validation of a novel flow-cytometric based method for the detection of inhibitory antibody responses. *J Immunol Methods* 2006;317(1–2):71–9.
91. Shamji MH, Ljorring C, Francis JN, et al. Functional rather than immunoreactive levels of IgG4 correlate closely with clinical response to grass pollen immunotherapy. *Allergy* 2012;67(2):217–26.
92. Feng M, Su Q, Lai X, et al. Functional and immunoreactive levels of IgG4 correlate with clinical responses during the maintenance phase of house dust mite immunotherapy. *J Immunol* 2018;200(12):3897–904.
93. Woo HY, Kim YS, Kang NI, et al. Mechanism for acute oral desensitization to antibiotics. *Allergy* 2006;61(8):954–8.
94. Novak N, Mete N, Bussmann C, et al. Early suppression of basophil activation during allergen-specific immunotherapy by histamine receptor 2. *J Allergy Clin Immunol* 2012;130:1153–8.
95. Schmid JM, Wurtzen PA, Dahl R, et al. Early improvement in basophil sensitivity predicts symptom relief with grass pollen immunotherapy. *J Allergy Clin Immunol* 2014;134(3):741–744 e745.
96. Shamji MH, Layhadi JA, Scadding GW, et al. Basophil expression of diamine oxidase: a novel biomarker of allergen immunotherapy response. *J Allergy Clin Immunol* 2015;135(4):913–21.e9.
97. Yuan X, Wang J, Li Y, et al. Allergy immunotherapy restores airway epithelial barrier dysfunction through suppressing IL-25 -induced endoplasmic reticulum stress in asthma. *Sci Rep* 2018;8(1):7950.
98. Valovirta E, Petersen TH, Piotrowska T, et al. Results from the 5-year SQ grass sublingual immunotherapy tablet asthma prevention (GAP) trial in children with grass pollen allergy. *J Allergy Clin Immunol* 2018;141(2):529–38.e13.
99. Dhimi S, Kakourou A, Asamoah F, et al. Allergen immunotherapy for allergic asthma: a systematic review and meta-analysis. *Allergy* 2017;72(12):1825–48.
100. Dhimi S, Nurmatov U, Arasi S, et al. Allergen immunotherapy for allergic rhinoconjunctivitis: a systematic review and meta-analysis. *Allergy* 2017;72(11):1597–631.
101. Tan HT, Hagner S, Ruchti F, et al. Tight junction, mucin, and inflammasome-related molecules are differentially expressed in eosinophilic, mixed, and neutrophilic experimental asthma in mice. *Allergy* 2019;74(2):294–307.
102. Russkamp D, Aguilar-Pimentel A, Alessandrini F, et al. IL-4 receptor alpha blockade prevents sensitization and alters acute and long-lasting effects of allergen-specific immunotherapy of murine allergic asthma. *Allergy* 2019;74(8):1549–60.